

# Rings of Bonds in Graphs

C. H. C. LITTLE AND D. A. HOLTON

*Department of Mathematics, Royal Melbourne Institute of Technology,  
124 la Trobe Street, Melbourne, Victoria 3000, Australia and  
Department of Mathematics, University of Melbourne, Parkville,  
Victoria 3052, Australia*

*Communicated by the Managing Editors*

Received July 15, 1977

It is shown that every strict maximal ring of bonds of size greater than 3 is even.

## 1. INTRODUCTION

Throughout we consider finite graphs. In addition, all graphs are assumed to be undirected unless specified otherwise. The reader is referred to Bondy and Murty [1] for any notation and terminology not explained here.

Let  $G$  be a graph and  $X, Y \subseteq V(G)$ . As in [1] we denote by  $[X, Y]$  the set of all edges joining a vertex of  $X$  and a vertex of  $Y$ . If  $\bar{X} = V(G) - X$ , then  $[X, \bar{X}]$  is called an *edge cut* when both  $X$  and  $\bar{X}$  are non-empty. We abbreviate  $[X, \bar{X}]$  by  $\delta(X)$  or  $\delta(\bar{X})$ . A minimal non-empty edge cut is a *bond*. If  $G$  is a directed graph and every edge is directed from the end in  $X$ , then  $\delta(X)$  is a *directed edge cut*.

Let  $R$  be a set of bonds. If the edges of  $G$  can be oriented so that every bond of  $R$  is directed, then we say that  $R$  is *consistently orientable*. A cyclic sequence of bonds  $R = (C_0, C_1, \dots, C_{n-1})$  with  $n \geq 3$  is a *ring* of bonds in the graph  $G$  if

- (i)  $R$  is consistently orientable.
- (ii)  $C_i \cap C_j \neq \emptyset$  if and only if  $i = j$ ,  $i \equiv j + 1 \pmod{n}$  or  $i \equiv j - 1 \pmod{n}$ , and
- (iii) no edge of  $G$  belongs to more than two bonds of  $R$ .

We note that (ii) implies (iii) except when  $n = 3$ .

For example, let  $G$  be the graph of Fig. 1. Then  $(\delta(\{v_1\}), \delta(\{v_2\}), \delta(\{v_3\}), \delta(\{v_4, v_5\}))$  is a ring, as is evident from the given orientations of edges, but  $(\delta(\{v_1\}), \delta(\{v_2\}), \delta(\{v_3\}), \delta(\{v_4\}), \delta(\{v_5\}))$  is not consistently orientable.

The *cardinality*,  $|R|$ , of  $R$  is the number of bonds in  $R$ . If  $R = (C_0,$




FIGURE 1

$C_1, \dots, C_{n-1}$ ), then  $R$  is referred to as an  $n$ -ring. A ring is *odd* or *even* according to whether  $|R|$  is odd or even.

The ring  $R$  is *strict* if there do not exist distinct bonds  $A, B, C$  satisfying the conditions  $B \in R$ ,  $C \in R$ ,  $B \cap C = \emptyset$ ,  $A \subseteq B \cup C$ . Furthermore,  $R$  is *maximal* if there does not exist a ring  $R' = (C'_0, C'_1, \dots, C'_{m-1})$  in  $G$  such that  $\bigcup_{k=0}^{m-1} C'_k \subseteq \bigcup_{l=0}^{n-1} C_l$  and  $m > n$ . We prove in this paper that every strict maximal ring with  $n \geq 4$  is even.

The dual of this result for planar graphs together with the main theorem of [3] proves a characterization of planar graphs conjectured in [4]. Chernyak [2] has independently also used the main theorem of [3] to prove this conjecture. Thus we have the following theorem, where the reader is referred to [3] for the relevant definitions.

**THEOREM 1.** *A graph is planar if and only if it has no maximal strict odd ring of circuits.*

## 2. $n$ -RINGS WITH $n \geq 4$

We begin by listing some elementary properties of bonds, each of which can be readily checked.

(1) Let  $G$  be a graph and let  $X$  be a non-empty proper subset of  $V(G)$ . The edge cut  $\delta(X)$  is a bond if and only if  $G[X]$  and  $G[\bar{X}]$  are connected. (See Exercise 2.2.8 of [1].)

(2) Let  $P = [S, \bar{S}]$  and  $Q = [T, \bar{T}]$  be bonds of a connected graph. Then

- (a)  $\delta(S \cap T) \subseteq P \cup Q$ ;
- (b) every bond from  $P \cup Q$  distinct from  $P$  and  $Q$  contains an edge of  $P - Q$  (and an edge of  $Q - P$ );
- (c)  $P \cap Q = [S \cap T, \bar{S} \cap \bar{T}] \cup [S \cap \bar{T}, \bar{S} \cap T]$ ;
- (d) if  $P$  and  $Q$  are directed bonds, then at least one of  $[S \cap T, \bar{S} \cap \bar{T}]$ ,  $[S \cap \bar{T}, \bar{S} \cap T]$  is empty.

Let  $R$  be a strict maximal  $n$ -ring  $(C_0, C_1, \dots, C_{n-1})$  of  $G$ , where  $n \geq 4$ . We henceforth assume that  $G$  is oriented so that every bond in  $R$  is directed. For each  $i \in \{0, 1, \dots, n-1\}$  we choose  $e_i \in C_i \cap C_{i+1}$ . (Throughout this paper, all subscripts are to be read modulo  $n$ .) Let  $C_i = \delta(A_i)$ . Since  $n \geq 4$  we have  $e_{i-2} \notin C_i$ . Therefore we may define  $D_i$  to be the element of  $\{A_i, \bar{A}_i\}$  which does not contain vertices incident with  $e_{i-2}$ . For all  $i$ , we define  $f(i) = 0$  if  $G$  is oriented so that every edge of  $C_i$  is directed toward the end in  $D_i$  and we let  $f(i) = 1$  otherwise.

LEMMA 2. *We have  $f(i) = f(i+1)$  if and only if*

$$C_i \cap C_{i+1} = [D_i \cap D_{i+1}, \bar{D}_i \cap \bar{D}_{i+1}].$$

*On the other hand,  $f(i) \neq f(i+1)$  if and only if*

$$C_i \cap C_{i+1} = [D_i \cap \bar{D}_{i+1}, \bar{D}_i \cap D_{i+1}].$$

COROLLARY 3. *If  $f(i) = f(i+1)$ , then  $[D_i \cap \bar{D}_{i+1}, \bar{D}_i \cap D_{i+1}] = \emptyset$ . If  $f(i) \neq f(i+1)$ , then  $[D_i \cap D_{i+1}, \bar{D}_i \cap \bar{D}_{i+1}] = \emptyset$ .*

This corollary follows as a result of (2(d)).

LEMMA 4. *If an edge of  $C_j$  joins two vertices of  $D_i$ , then  $j \equiv i \pm 1 \pmod{n}$ .*

*Proof.* Suppose that  $e$  is an edge of  $C_j$  that joins two vertices of  $D_i$ , where  $j \not\equiv i \pm 1 \pmod{n}$ .

Suppose further that for every  $k \notin \{i-1, i, i+1\}$ , either  $C_k \cap [D_i, D_i] = \emptyset$  or  $C_k \cap [\bar{D}_i, \bar{D}_i] = \emptyset$ . Recall that  $e_{i-2}$  joins two vertices of  $\bar{D}_i$ . Hence every edge of  $C_{i-2}$  joins vertices of  $\bar{D}_i$  by assumption. Since  $C_{i-2} \cap C_{i-3} \neq \emptyset$ , the same is also true of  $C_{i-3}$  (if  $n > 4$ ) and, by induction, of  $C_j$ , in contradiction to the existence of  $e$ . We conclude that for some  $k \notin \{i-1, i, i+1\}$ ,  $C_k \cap [D_i, D_i] \neq \emptyset$  and  $C_k \cap [\bar{D}_i, \bar{D}_i] \neq \emptyset$ .

By (2(a)),  $\delta(D_i \cap D_k) \subseteq C_i \cup C_k$ . Hence some subset of  $\delta(D_i \cap D_k)$  is a bond  $C$  such that  $C \subseteq C_i \cup C_k$ . Since  $C_k \cap [\bar{D}_i, \bar{D}_i] \neq \emptyset$  we have  $C \neq C_k$ . Suppose  $C_i \cap [\bar{D}_k, \bar{D}_k] = \emptyset$ . Then  $[D_i \cap \bar{D}_k, \bar{D}_i \cap \bar{D}_k] = \emptyset$ , so that  $\delta(D_i \cap \bar{D}_k) = [D_i \cap \bar{D}_k, \bar{D}_i \cap \bar{D}_k] \subseteq C_k$  since  $C_k \cap [\bar{D}_i, \bar{D}_i] \neq \emptyset$ . On the other hand,  $\delta(D_i \cap \bar{D}_k) \neq \emptyset$  because  $D_i \cap \bar{D}_k \neq \emptyset$  and  $G$  is connected. These results contradict the fact that  $C_k$  is a bond. Hence  $C_i \cap [\bar{D}_k, \bar{D}_k] \neq \emptyset$ , and so  $C \neq C_i$ . Since  $k \notin \{i-1, i, i+1\}$  we have  $C_i \cap C_k = \emptyset$  and thus the strictness of  $R$  is contradicted. ■

COROLLARY 5.  *$C_j \subseteq [\bar{D}_i, \bar{D}_i]$  for all  $j \notin \{i-1, i, i+1\}$ .*

LEMMA 6. *For each  $i$ ,  $f(i) \neq f(i+1)$ .*

*Proof.* Assume  $f(i) = f(i+1)$  for some  $i$ , and without loss of generality, take  $f(i) = 0$ . We shall contradict the maximality of  $R$  by constructing a ring  $\hat{R}$  in which  $C_i$  and  $C_{i+1}$  are replaced by three bonds:  $\hat{C}_i$ ,  $\hat{C}$  and  $\hat{C}_{i+1}$ .

Since  $e_{i-1} \in C_i = [D_i, \bar{D}_i]$  and by Corollary 5,  $e_{i-1} \in C_{i-1} \subseteq [\bar{D}_{i+1}, \bar{D}_{i+1}]$ , then  $e_{i-1} \in [D_i \cap \bar{D}_{i+1}, \bar{D}_i \cap \bar{D}_{i+1}]$ . Let  $\hat{D}_i$  be the vertex set of the component of  $G[D_i \cap \bar{D}_{i+1}]$  containing an end of  $e_{i-1}$ .

We show that  $\hat{C}_i = \delta(\hat{D}_i)$  is a bond. Let  $\hat{G} = G[V(G) - \hat{D}_i]$ . If  $L$  is a component of  $\hat{G}$  containing vertices of  $D_{i+1}$ , then  $L$  contains all vertices of  $D_{i+1}$  since  $G[D_{i+1}]$  is connected and  $\hat{D}_i \subseteq \bar{D}_{i+1}$ . Now  $[D_i \cap D_{i+1}, \bar{D}_i] \neq \emptyset$  by Lemma 2, since otherwise  $C_{i+1} \cap C_i = \emptyset$ . Thus  $G[\bar{D}_i]$  is a subgraph of  $L$ . Therefore any component  $M$  of  $\hat{G}$  other than  $L$  contains only vertices of  $D_i \cap \bar{D}_{i+1}$ . Since  $G$  is connected, we have  $[V(M), \hat{D}_i] \neq \emptyset$ , so that the existence of  $M$  contradicts the definition of  $\hat{D}_i$ . Thus  $L$  is the only component of  $\hat{G}$  and it follows that  $\hat{C}_i$  is a bond.

Similarly, for any  $X$  such that  $G[X]$  is a component of  $G[\bar{D}_i \cap D_{i+1}]$ ,  $\delta(X)$  is a bond. Let  $\mathcal{D}$  be the union of the vertex sets of all such  $X$  for which  $\delta(X) \cap C_{i+2} \neq \emptyset$ .

By Lemma 2,  $C_i \cap C_{i+1} = [D_i \cap D_{i+1}, \bar{D}_i \cap \bar{D}_{i+1}]$ . Hence  $e_i$  has neither end in  $D_i \cap \bar{D}_{i+1}$  and so  $\hat{C}_i \subseteq (C_i \cap C_{i+1}) - \{e_i\}$ . Thus  $\hat{C}_i$  is distinct from  $C_i$  and  $C_{i+1}$  which implies that  $\hat{C}_i$  contains an edge  $\hat{e}_i \in C_{i+1} - C_i$ . In particular,  $\hat{e}_i \in [D_i \cap \bar{D}_{i+1}, D_i \cap D_{i+1}]$ . Thus there is a component of  $G[D_{i+1} \cap \mathcal{D}]$  containing an end of  $\hat{e}_i$ . Call the vertex set of this component  $\hat{D}$ .

We now show that  $\hat{C} = \delta(\hat{D})$  is a bond. Since  $\hat{D} \subseteq D_{i+1}$  and  $C_{i+1}$  is a bond,  $\bar{D}_{i+1}$  is contained in a single component  $Y$  of  $G - \hat{D}$ . Choose a component  $Z$  of  $G - \hat{D}$ . It suffices to show that  $Z = Y$ . If  $V(Z) \cap \mathcal{D} \neq \emptyset$ , then for some vertex set  $X$  such that  $G[X]$  is a component of  $G[\mathcal{D}]$ , we have  $X \subseteq V(Z)$ . Because  $G[X]$  is a component of  $G[\bar{D}_i \cap D_{i+1}]$ , any edge  $e$  of  $\delta(X)$  must have an end  $w \in D_i \cup \bar{D}_{i+1}$ . If  $w \in \bar{D}_{i+1}$ , then  $Z = Y$ . Hence we assume that  $\delta(X) = [X, D_i]$ . But now we have the contradiction that  $\delta(X) \cap C_{i+2} = \emptyset$  by Corollary 5. On the other hand, suppose  $V(Z) \cap \mathcal{D} = \emptyset$ . If  $Z \neq Y$ , then  $V(Z) \subseteq D_{i+1} \cap \mathcal{D}$ . Thus  $G[\hat{D}]$  cannot be a component of  $G[D_{i+1} \cap \mathcal{D}]$  because  $[\hat{D}, V(Z)] \neq \emptyset$  by the fact that  $G$  is connected. Therefore  $Z = Y$ , so that  $\hat{C}$  is a bond.

Suppose  $e_{i+1} \in \hat{C}$ . Then  $e_{i+1}$  joins a vertex  $x \in D_{i+1} \cap \mathcal{D}$  to a vertex  $y \in \bar{D}_{i+1}$ , since  $e_{i+1} \in C_{i+1}$ . But  $e_{i+1} \in C_{i+2}$ , and so  $x \in \bar{D}_i$  by Corollary 5. Hence  $e \in \delta(\bar{D}_i \cap D_{i+1})$  and since  $e_{i+1} \in C_{i+2}$  the fact that  $x \in \mathcal{D}$  is contradicted. Thus  $\hat{C} \subseteq C_i \cup C_{i+1} - \{e_{i-1}, e_{i+1}\}$ . Hence  $\hat{C}$  contains an edge  $\hat{e} \in C_i - C_{i+1}$  by (2(b)). Then  $\hat{e} \in [D_{i+1} \cap \mathcal{D}, \mathcal{D}]$ . Now we define  $\hat{D}_{i+1}$  to be the vertex set of the component of  $G[\mathcal{D}]$  containing an end of  $\hat{e}$ . Let  $\hat{C}_{i+1} = \delta(\hat{D}_{i+1})$ . We have already shown that  $\hat{C}_{i+1}$  is a bond because  $G[\bar{D}_{i+1}]$  is a component of  $G[\mathcal{D}]$ .

Define  $\hat{R} = (C_0, C_1, \dots, C_{i-1}, \hat{C}_i, \hat{C}, \hat{C}_{i+1}, C_{i+2}, \dots, C_{n-1})$ . We now aim to

show that  $\hat{R}$  is ring of bonds and thus contradict the maximality of  $R$ . First we reverse the direction of every edge of  $\hat{C} \cup \{([D_i, D_i] \cup C_i) \cap ([D_{i+1}, D_{i+1}] \cup C_{i+1})\}$ . Since  $f(i) = 0$  each edge of  $C_i$  was directed toward the end in  $D_i$  in the original orientation of  $G$ .

Now we show that  $\hat{R}$  is consistently orientable.

First we recall that  $\hat{C} \subseteq \delta(D_{i+1} \cap \mathcal{D})$ , so that by Corollary 5 and the definition of a ring of bonds,  $\hat{C}$  contains no edge of any bond in  $R - \{C_i, C_{i+1}, C_{i+2}\}$ . Hence it is clear that these bonds remain directed under the new orientation of  $G$ .

We have  $\hat{C} \cap C_{i+2} = \emptyset$ , for otherwise there is an edge of  $\hat{C} \cap C_{i+2}$  in  $[\bar{D}_i, \bar{D}_i]$ , in contradiction to the definition of  $\mathcal{D}$ . Furthermore  $C_{i+2} \cap [D_i, D_i] = \emptyset$  and  $C_{i+2} \cap C_i = \emptyset$  so that  $C_{i+2}$  is a directed bond under the new orientation of  $G$ .

Now  $\hat{C}_i \subseteq (C_i \cap [\bar{D}_{i+1}, \bar{D}_{i+1}]) \cup (C_{i+1} \cap [D_i, D_i])$  by Corollary 3. The edges of  $C_i \cap [\bar{D}_{i+1}, \bar{D}_{i+1}]$  are directed towards  $\hat{D}_i$  both before and after the reorientation of  $G$ . When  $G$  is reoriented the edges of  $C_{i+1} \cap [D_i, D_i]$  are also directed towards  $\hat{D}_i$ . Hence  $\hat{C}_i$  is a directed bond in the new orientation of  $G$ .

Next consider  $\hat{C}$ . Every edge in  $\hat{C}$  has an end in  $\hat{D} \cap D_{i+1}$  and every edge in  $\hat{C} - C_{i+1}$  has an end in  $\hat{D} \cap D_i$  by the definition of  $\mathcal{D}$ . Thus  $\hat{C}$  is directed in both the old and new orientations of  $G$ .

In considering  $\hat{C}_{i+1}$  we note that if there exists an edge  $e' \in \hat{C}_{i+1} \cap [D_i, \hat{D}_{i+1}]$ , then  $e' \in [\bar{D}_i \cap \bar{D}_{i+1}, \hat{D}_{i+1} \cap \bar{D}_i \cap D_{i+1}]$ , and hence is directed towards  $\hat{D}_{i+1}$  in the new orientation. Hence  $\hat{C}_{i+1}$  is a directed bond. Thus  $\hat{R}$  is consistently orientable.

By construction we note that  $e_{i-1} \in C_{i-1} \cap \hat{C}_i$ ,  $\hat{e}_i \in \hat{C}_i \cap \hat{C}$ ,  $\hat{e} \in \hat{C} \cap \hat{C}_{i+1}$  and  $\hat{C}_{i+1} \cap C_{i+2} \neq \emptyset$ . Hence consecutive bonds of  $\hat{R}$  are not disjoint.

It remains to show that non-consecutive pairs of bonds in  $\hat{R}$  are disjoint. Since  $\hat{C}_i$ ,  $\hat{C}$  and  $\hat{C}_{i+1}$  are all subsets of  $C_i \cup C_{i+1}$ , they are disjoint from  $C_{i+3}, C_{i+4}, \dots, C_{i-2}$ . This leaves only the five cases below to be considered.

- (i)  $C_{i-1} \cap \hat{C} = \emptyset$  (as has already been noted) since  $C_{i-1} \subseteq [\bar{D}_{i+1}, \bar{D}_{i+1}]$  by Corollary 5.
- (ii)  $C_{i-1} \cap \hat{C}_{i+1} = \emptyset$  by the same argument.
- (iii)  $\hat{C}_i \cap \hat{C}_{i+1} = \emptyset$  by Corollary 3.
- (iv)  $\hat{C}_i \cap C_{i+2} = \emptyset$  since  $C_{i+2} \subseteq [\bar{D}_i, \bar{D}_i]$  by Corollary 5.
- (v)  $\hat{C} \cap C_{i+2} = \emptyset$  as has already been shown.

Hence  $\hat{R}$  is a ring of bonds. This conclusion contradicts the maximality of  $R$  and hence  $f(i) \neq f(i+1)$  for each  $i$ . ■

The main result of this paper is immediate.

**THEOREM 7.** *If  $n \geq 4$ , then every strict maximal  $n$ -ring of bonds is even.*

## ACKNOWLEDGMENT

We are deeply indebted to the referees of earlier versions of this work, from the result of whose considerable effort this much simpler paper has evolved.

## REFERENCES

1. J. A. BONDY AND U. S. R. MURTY, "Graph Theory with Applications," Macmillan & Co., London, 1976.
2. A. A. CHERNYAK, The Little hypothesis on planar graphs, *Izv. Akad. Nauk BSSR Ser. Fiz. Mat.* **2** (1980), 41–45. [In Russian.]
3. D. A. HOLTON AND C. H. C. LITTLE, The non-planarity of maximal 3-rings of circuits, submitted for publication.
4. C. H. C. LITTLE, On rings of circuits in planar graphs, in "Proceedings, 5th Austral. Conf. Comb. Math.," pp. 133–140, Lecture Notes in Mathematics No. 622, Springer-Verlag, Berlin, 1977.